Rangeland Productivity and Plant Composition

United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

| Map symbol and soil name | Ecological site | Total | lry-weight prod | uction | Characteristic vegetation | Rangeland |
|-----------------------------|-----------------|-------------------|-----------------|---------------------|---------------------------|-------------|
| | | Favorable year | Normal year | Unfavorable year | | composition |
| | | Lb/ac | Lb/ac | Lb/ac | | Pa |
| s1853: | | | | | | |
| Portneuf | | | | | | |
| Trevino | | 700 | 500 | 250 | Bluebunch wheatgrass | 2 |
| | | | | | Wyoming big sagebrush | 2 |
| | | | | | Other perennial forbs | 1(|
| | | | | | Sandberg bluegrass | 1(|
| | | | | | Thurber needlegrass | 1(|
| | | | | | Arrowleaf balsamroot | Ę |
| | | | | | Longleaf hawksbeard | Ę |
| | | | | | Other perennial grasses | : |
| | | | | | Other shrubs | ţ |
| Trevino | | 700 | 500 | 250 | Bluebunch wheatgrass | 25 |
| | | | | | Wyoming big sagebrush | 25 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Sandberg bluegrass | 10 |
| | | | | | Thurber needlegrass | 10 |
| | | | | | Arrowleaf balsamroot | Ę |
| | | | | | Longleaf hawksbeard | ţ |
| | | | | | Other perennial grasses | Ę |
| | | | | | Other shrubs | Ę |



Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total o | lry-weight prod | uction | Characteristic vegetation | Rangeland composition |
|-----------------------------|-------------------------|-------------------|-----------------|---------------------|---------------------------|-----------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s1853: | | | | | | |
| Taunton | | 1,100 | 700 | 300 | Bluebunch wheatgrass | 30 |
| | | | | | Wyoming big sagebrush | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Sandberg bluegrass | 10 |
| | | | | | Longleaf hawksbeard | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Phlox | 5 |
| Portneuf | LOAMY 8-12 ARTRW8/PSSP6 | 1,200 | 700 | 500 | Bluebunch wheatgrass | 30 |
| | | | | | Wyoming big sagebrush | 25 |
| | | | | | Thurber needlegrass | 20 |
| | | | | | Green rabbitbrush | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Prairie Junegrass | 5 |
| | | | | | Sandberg bluegrass | 5 |
| | | | | | Threetip sagebrush | 5 |
| Rock outcrop | | | | | | |
| Scoon | | 500 | 350 | 200 | Bluebunch wheatgrass | 65 |
| | | | | | Sandberg bluegrass | 10 |
| | | | | | Big sagebrush | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Thurber needlegrass | 5 |
| | | | | | Buckwheat | 3 |
| | | | | | Balsamroot | 2 |
| | | | | | Biscuitroot | 1 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland |
|-----------------------------|-------------------------|-----------------------------|----------------|---------------------|---------------------------|-------------|
| | | Favorable year | Normal year | Unfavorable year | | composition |
| | • | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s1853: | | | | | | |
| Declo | LOAMY 8-12 ARTRW8/PSSP6 | 1,100 | 700 | 500 | Bluebunch wheatgrass | 30 |
| | | | | | Other perennial grasses | 15 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Wyoming big sagebrush | 15 |
| | | | | | Arrowleaf balsamroot | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Rabbitbrush | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Wodskow | | | | | | |

