United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland |
|--------------------------|-----------------------------|-----------------------------|----------------|------------------|---------------------------|-------------|
| | | Favorable year | Normal year | Unfavorable year | Characteristic vegetation | composition |
| | | Lb/ac | Lb/ac | Lb/ac | | Pc |
| s1981: | | | | | | |
| Royal | SANDY LOAM 8-12 ARTRW8/ACHY | 900 | 600 | 350 | Wyoming big sagebrush | 30 |
| • | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Phlox | 5 |
| | | | | | Sand dropseed | 5 |
| | | | | | Sandberg bluegrass | 5 |
| | | | | | Spiny hopsage | 5 |
| Xeric Torriorthents | SANDY LOAM 8-12 ARTRW8/ACHY | 900 | 600 | 350 | Wyoming big sagebrush | 30 |
| | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Bottlebrush squirreltail | 7 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Sandberg bluegrass | 5 |
| | | | | | Arrowleaf balsamroot | 3 |
| Quincy | SAND 8-12 ARTRT/ACHY | 950 | 650 | 450 | Basin big sagebrush | 30 |
| | | | | | Indian ricegrass | 25 |
| | | | | | Needle and thread | 15 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Sand dropseed | 5 |



| Map symbol and soil name | Ecological site | Total | dry-weight prod | uction | Characteristic vegetation | Rangeland composition |
|--------------------------|-----------------------------|-------------------|-----------------|------------------|---------------------------|-----------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | Lb/ac | Lb/ac | Lb/ac | | Pc |
| s1981: | | | | | | |
| Xerollic Camborthids | SANDY LOAM 8-12 ARTRW8/ACHY | 1,000 | 600 | 350 | Wyoming big sagebrush | 35 |
| | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Other shrubs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Davey | SANDY LOAM 8-12 ARTRW8/ACHY | 1,000 | 600 | 350 | Wyoming big sagebrush | 35 |
| • | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Other shrubs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Royal | SANDY LOAM 8-12 ARTRW8/ACHY | 900 | 600 | 350 | Wyoming big sagebrush | 30 |
| • | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Phlox | 5 |
| | | | | | Sand dropseed | 5 |
| | | | | | Sandberg bluegrass | 5 |
| | | | | | Spiny hopsage | 5 |



| Map symbol and soil name | Ecological site | Total o | lry-weight prod | uction | Characteristic vegetation | Rangeland composition |
|--------------------------|-----------------------------|-------------------|-----------------|------------------|---------------------------|-----------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s1981: | | | | | | |
| Truesdale | SANDY LOAM 8-12 ARTRW8/ACHY | 1,000 | 600 | 350 | Wyoming big sagebrush | 35 |
| | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Other shrubs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Shano | SANDY LOAM 8-12 ARTRW8/ACHY | 1,000 | 600 | 350 | Wyoming big sagebrush | 35 |
| | | · | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Other shrubs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Bahem | LOAMY 7-10 ARTRW8/STTH2 | 700 | 500 | 300 | Wyoming big sagebrush | 35 |
| | | | | | Thurber needlegrass | 30 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Rabbitbrush | 5 |
| | | | | | Sandberg bluegrass | 5 |



| Map symbol and soil name | Ecological site | Total | dry-weight prod | uction | Characteristic vegetation | Rangeland composition |
|--------------------------|-----------------------------|-------------------|-----------------|------------------|---------------------------|-----------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | Lb/ac | Lb/ac | Lb/ac | | Pc |
| s1981: | | | | | | |
| Buko | SANDY LOAM 8-12 ARTRW8/ACHY | 1,000 | 600 | 350 | Wyoming big sagebrush | 35 |
| | | | | | Indian ricegrass | 20 |
| | | | | | Thurber needlegrass | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Other shrubs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Davey | SAND 8-12 ARTRT/ACHY | 950 | 650 | 450 | Basin big sagebrush | 30 |
| | | | | | Indian ricegrass | 25 |
| | | | | | Needle and thread | 15 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Sand dropseed | 5 |
| Purdam | LOAMY 7-10 ARTRW8/STTH2 | 700 | 500 | 300 | Wyoming big sagebrush | 35 |
| | | | | | Thurber needlegrass | 30 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Rabbitbrush | 5 |
| | | | | | Sandberg bluegrass | 5 |
| Dune land | | | | | | |



| Map symbol | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland |
|---------------|-------------------------|-----------------------------|----------------|------------------|---------------------------|-------------|
| and soil name | | Favorable year | Normal year | Unfavorable year | Characteriotic vogetation | composition |
| | | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s1981: | | | | | | |
| Elijah | LOAMY 7-10 ARTRW8/STTH2 | 700 | 500 | 300 | Wyoming big sagebrush | 35 |
| | | | | | Thurber needlegrass | 30 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Rabbitbrush | 5 |
| | | | | | Sandberg bluegrass | 5 |

