United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland |
|-----------------------------|-------------------------|-----------------------------|----------------|---------------------|---------------------------|-------------|
| | | Favorable year | Normal year | Unfavorable year | onaracionsile vegetation | composition |
| | • | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s2157: | | | | | | |
| Lanoak | | | | | | |
| Manila | | | | | | |
| | | | | | | |
| Brifox | LOAMY 13-16 ARTRV/PSSP6 | 1,600 | 1,300 | 900 | Bluebunch wheatgrass | 25 |
| | | | | | Mountain big sagebrush | 20 |
| | | | | | Other perennial grasses | 15 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Prairie junegrass | 10 |
| | | | | | Common snowberry | 5 |
| | | | | | Geranium | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Rabbitbrush | 5 |
| Huffman | | | | | | |
| Bear Lake | WET MEADOW | 4,500 | 3,600 | 3,000 | Tufted hairgrass | 20 |
| | | | | | Cinquefoil | 5 |
| | | | | | Clover | 5 |
| | | | | | Kentucky bluegrass | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| Broadhead | | | | | | |
| Huffman | | | | | | |



United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland |
|-----------------------------|-------------------------------|-----------------------------|----------------|---------------------|---------------------------|-------------|
| | | Favorable year | Normal year | Unfavorable year | | composition |
| | | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s2157: | | | | | | |
| Oxford | STEEP SLOPE 12-16 ARTRT/PSSP6 | 1,800 | 1,200 | 800 | Bluebunch wheatgrass | 35 |
| | | | | | Basin big sagebrush | 15 |
| | | | | | Antelope bitterbrush | 5 |
| | | | | | Arrowleaf balsamroot | 5 |
| | | | | | Western wheatgrass | 5 |
| Bergquist | MOUNTAIN LOAM 18-22 | 2,100 | 1,900 | 1,600 | Bigtooth maple | 10 |
| | ACSAG2/ARTRV/PSSP6 | , | , | , | Bluebunch wheatgrass | 10 |
| | | | | | Mountain big sagebrush | 10 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Other perennial grasses | 10 |
| | | | | | Other shrubs | 10 |
| | | | | | Common chokecherry | 5 |
| | | | | | Nevada bluegrass | 5 |
| | | | | | Snowberry | 5 |
| Holmes | RIVERBOTTOM | 2,000 | 1,500 | 1,200 | Cottonwood | 15 |
| | | | | | Willow | 15 |
| | | | | | Bluebunch wheatgrass | 10 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |



United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|---------------------------------|-----------------------------|----------------|---------------------|--|-----------------------|
| | | Favorable year | Normal year | Unfavorable year | - | composition |
| | • | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s2157: | | | | | | |
| Merkley | LOAMY BOTTOM 12-16 ARTRT/LECI4- | 1,800 | 1,500 | 1,200 | Basin big sagebrush | 20 |
| | AGROP | | | | Thickspike wheatgrass | 15 |
| | | | | | Bluebunch wheatgrass | 10 |
| | | | | | Arrowleaf balsamroot | 5 |
| | | | | | Green rabbitbrush | 5 |
| | | | | | Longleaf hawksbeard Other shrubs | 5 5 |
| | | | | | Western wheatgrass | 5 |
| | | | | | Western Wheatgrass | 5 |
| Xerorthents | STEEP SLOPES 12-16 ARTRV/PSSP6 | 1,500 | 1,100 | 550 | Bluebunch wheatgrass | 20 |
| | | | | | Mountain big sagebrush | 20 |
| | | | | | Longleaf hawksbeard | 5 |
| | | | | | Nevada bluegrass | 5 |
| | | | | | Sticky geranium | 5 |
| | | | | | Thickspike wheatgrass | 5 |
| Banida | STEEP SLOPE 12-16 ARTRT/PSSP6 | 1,600 | 1,000 | 800 | Bluebunch wheatgrass | 35 |
| Danida | | 1,000 | 1,000 | 000 | Basin big sagebrush | 20 |
| | | | | | Antelope bitterbrush | 5 |
| | | | | | Arrowleaf balsamroot | 5 |
| | | | | | Western wheatgrass | 5 |
| Lana | SEMIWET MEADOW | 0.400 | 0.000 | 0.000 | | 00 |
| Lago | SEMIWET MEADOW | 3,400 | 2,600 | 2,000 | Slender wheatgrass Tufted hairgrass | 20 15 |
| | | | | | Baltic rush | 5 |
| | | | | | Cinquefoil | 5 |
| | | | | | • | 5 |
| | | | | | Cinquetoil Fowl bluegrass | |



United States

| Map symbol and soil name | Ecological site | Total o | lry-weight prod | uction | Characteristic vegetation | Rangeland composition |
|-----------------------------|---------------------------------|-------------------|-----------------|---------------------|---------------------------|-----------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | · | Lb/ac | Lb/ac | Lb/ac | | Pct |
| s2157: | | | | | | |
| Lanoak | LOAMY 16-22 ARTRV/PSSP6 16-22" | 2,500 | 1,600 | 1,000 | Bluebunch wheatgrass | 30 |
| | | | | | Idaho fescue | 10 |
| | | | | | Mountain big sagebrush | 10 |
| | | | | | Antelope bitterbrush | 5 |
| | | | | | Arrowleaf balsamroot | 5 |
| | | | | | Columbia needlegrass | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Prairie junegrass | 5 |
| | | | | | Sticky geranium | 5 |
| | | | | | Western snowberry | 5 |
| Vitale | GRAVELLY LOAM 16-22 ARTRV/PSSP6 | 2,000 | 1,500 | 850 | Bluebunch wheatgrass | 25 |
| | | | | | Mountain big sagebrush | 15 |
| | | | | | Arrowleaf balsamroot | 5 |
| | | | | | Longleaf hawksbeard | 5 |
| | | | | | Lupine | 5 |
| | | | | | Nevada bluegrass | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| Yeates Hollow | STONY LOAM 16-22 ARTRV/PSSP6 | 1,800 | 1,100 | 600 | Bluebunch wheatgrass | 40 |
| | | | , | | Antelope bitterbrush | 5 |
| | | | | | Columbia needlegrass | 5 |
| | | | | | Cutleaf balsamroot | 5 |
| | | | | | Geranium | 5 |
| | | | | | Mountain big sagebrush | 5 |
| | | | | | Slender wheatgrass | 5 |

