

Rangeland Productivity and Plant Composition

United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland composition
		Favorable year	Normal year	Unfavorable year		
		<i>Lb/ac</i>	<i>Lb/ac</i>	<i>Lb/ac</i>		<i>Pct</i>
s5565:						
Ocala	---	2,000	1,000	500	Alkali sacaton Basin wildrye Black greasewood Basin big sagebrush Inland saltgrass Rubber rabbitbrush	25 15 10 5 5 5
Ocala	---	1,900	1,400	800	Basin wildrye Alkali sacaton Black greasewood Inland saltgrass Rubber rabbitbrush	50 20 10 10 5
Valmy	---	800	600	350	Black greasewood Basin big sagebrush Spiny hopsage Basin wildrye Bottlebrush squirreltail Indian ricegrass	30 25 15 10 5 5
Sondoa	---	600	450	200	Black greasewood Alkali sacaton Inland saltgrass	55 10 5
Kawich	---	700	500	350	Indian ricegrass Black greasewood Fourwing saltbush Bottlebrush squirreltail Rubber rabbitbrush	30 15 15 5 5

Rangeland Productivity and Plant Composition

United States

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland composition
		Favorable year	Normal year	Unfavorable year		
		<i>Lb/ac</i>	<i>Lb/ac</i>	<i>Lb/ac</i>		<i>Pct</i>
s5565:						
Playas	---	---	---	---	---	---
Sonoma	---	1,500	1,000	700	Alkali sacaton Alkali bluegrass Alkali cordgrass Alkali muhly Inland saltgrass Basin wildrye	30 10 10 10 10 5
Water	---	---	---	---	---	---