

Rangeland Productivity and Plant Composition

United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|--|---|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5666: Zadvar | --- | 700 | 500 | 250 | Black sagebrush Galleta Indian ricegrass Winterfat | 30 10 10 5 |
| Veet | --- | 900 | 600 | 300 | Indian ricegrass Wyoming big sagebrush Spiny hopsage Bottlebrush squirreltail Bud sagebrush Desert needlegrass Galleta Needleandthread Nevada ephedra Winterfat | 20 20 10 5 5 5 5 5 5 5 |
| Handpah | --- | 800 | 500 | 300 | Wyoming big sagebrush Galleta Indian ricegrass Bottlebrush squirreltail Desert needlegrass Fourwing saltbush Spiny hopsage | 20 10 10 5 5 5 5 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | |
| s5666: | | | | | | <i>Pct</i> |
| Handpah | --- | 800 | 500 | 300 | Wyoming big sagebrush | 20 |
| | | | | | Galleta | 10 |
| | | | | | Indian ricegrass | 10 |
| | | | | | Black sagebrush | 5 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Desert needlegrass | 5 |
| | | | | | Douglas rabbitbrush | 5 |
| | | | | | Fourwing saltbush | 5 |
| | | | | | Spiny hopsage | 5 |
| Armespan | --- | 700 | 400 | 200 | Black sagebrush | 25 |
| | | | | | Galleta | 10 |
| | | | | | Indian ricegrass | 10 |
| | | | | | Bud sagebrush | 5 |
| | | | | | Nevada ephedra | 5 |
| | | | | | Winterfat | 5 |
| | | | | | Bottlebrush squirreltail | 2 |
| | | | | | Needlegrass | 1 |
| Breko | --- | 800 | 500 | 300 | Wyoming big sagebrush | 20 |
| | | | | | Galleta | 10 |
| | | | | | Desert needlegrass | 5 |
| | | | | | Fourwing saltbush | 5 |
| | | | | | Indian ricegrass | 5 |
| | | | | | Nevada ephedra | 5 |
| | | | | | Spiny hopsage | 5 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5666: | | | | | | |
| Handpah | --- | 800 | 500 | 300 | Wyoming big sagebrush | 20 |
| | | | | | Galleta | 10 |
| | | | | | Indian ricegrass | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Desert needlegrass | 5 |
| | | | | | Fourwing saltbush | 5 |
| | | | | | Spiny hopsage | 5 |
| Wrango | --- | 900 | 700 | 400 | Black sagebrush | 25 |
| | | | | | Indian ricegrass | 20 |
| | | | | | Basin wildrye | 5 |
| | | | | | Bud sagebrush | 5 |
| | | | | | Needleandthread | 5 |
| | | | | | Winterfat | 5 |
| | | | | | Galleta | 2 |
| Armespan variant | --- | --- | --- | --- | --- | --- |
| Lyda | --- | 400 | 200 | 100 | Bailey greasewood | 15 |
| | | | | | Spiny menodora | 15 |
| | | | | | Bud sagebrush | 10 |
| | | | | | Galleta | 10 |
| | | | | | Indian ricegrass | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Cooper wolfberry | 5 |
| | | | | | Desert needlegrass | 5 |
| | | | | | Nevada ephedra | 5 |
| | | | | | Shadscale | 5 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|--|---|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5666: Unsel | --- | 350 | 250 | 100 | Shadscale Galleta Bud sagebrush Indian ricegrass Winterfat Bailey greasewood Bottlebrush squirreltail Desert needlegrass Fourwing saltbush Nevada ephedra | 20 15 10 10 10 5 5 5 5 5 |
| Ubehebe | --- | --- | --- | --- | --- | --- |
| Veet variant | --- | 3,000 | 2,000 | 800 | Basin wildrye Basin big sagebrush Fourwing saltbush Rubber rabbitbrush Western wheatgrass Wyoming big sagebrush | 25 5 5 5 5 5 |