

Rangeland Productivity and Plant Composition

United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|--------------------------|-----------------|-----------------------------|--------------|------------------|--|------------------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5853: | | | | | | |
| Ocala | --- | 500 | 350 | 200 | Black greasewood Basin wildrye Inland saltgrass | 60 10 5 |
| Ocala | --- | 1,900 | 1,400 | 800 | Basin wildrye Alkali sacaton Black greasewood Inland saltgrass Rubber rabbitbrush | 50 20 10 10 5 |
| Humboldt | --- | 3,000 | 2,500 | 1,800 | Basin wildrye Creeping wildrye Willow Alkali sacaton Basin big sagebrush Inland saltgrass Nevada bluegrass | 30 25 10 5 5 5 5 |
| Humboldt | --- | 3,000 | 2,500 | 1,800 | Basin wildrye Creeping wildrye Willow Alkali sacaton Basin big sagebrush Inland saltgrass Nevada bluegrass | 30 25 10 5 5 5 5 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5853: | | | | | | |
| Humboldt | --- | 1,900 | 1,400 | 800 | Basin wildrye | 50 |
| | | | | | Alkali sacaton | 20 |
| | | | | | Black greasewood | 10 |
| | | | | | Inland saltgrass | 5 |
| | | | | | Rabbitbrush | 5 |
| Griver | --- | 3,000 | 2,400 | 1,800 | Basin wildrye | 30 |
| | | | | | Creeping wildrye | 30 |
| | | | | | Other perennial grasses | 7 |
| | | | | | Coyote willow | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Sedge | 5 |
| | | | | | Woods rose | 5 |
| | | | | | Baltic rush | 3 |
| | | | | | Cinquefoil | 2 |
| | | | | | Groundsel | 2 |
| | | | | | Other perennial forbs | 2 |
| | | | | | Wildiris | 2 |
| | | | | | Yarrow | 2 |
| Iron Blossom | --- | 600 | 450 | 300 | Shadscale | 40 |
| | | | | | Black greasewood | 25 |
| | | | | | Bottlebrush squirreltail | 10 |
| | | | | | Bud sagebrush | 5 |
| | | | | | Seepweed | 5 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5853: | | | | | | |
| Iron Blossom | --- | 1,500 | 1,000 | 600 | Basin wildrye | 50 |
| | | | | | Basin big sagebrush | 15 |
| | | | | | Black greasewood | 10 |
| | | | | | Bottlebrush squirreltail | 5 |
| | | | | | Povertyweed | 5 |
| | | | | | Rubber rabbitbrush | 5 |
| | | | | | Western wheatgrass | 5 |
| Rixie | --- | 1,900 | 1,400 | 800 | Basin wildrye | 40 |
| | | | | | Alkali sacaton | 15 |
| | | | | | Black greasewood | 10 |
| | | | | | Inland saltgrass | 10 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Povertyweed | 5 |
| | | | | | Rubber rabbitbrush | 5 |
| Griver | --- | 2,500 | 1,800 | 1,200 | Basin wildrye | 50 |
| | | | | | Big sagebrush | 10 |
| | | | | | Lupine | 5 |
| | | | | | Mat muhly | 5 |
| | | | | | Nevada bluegrass | 5 |
| | | | | | Other perennial forbs | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Rubber rabbitbrush | 5 |
| | | | | | Western wheatgrass | 5 |
| | | | | | Bottlebrush squirreltail | 2 |
| | | | | | Sandberg bluegrass | 2 |
| | | | | | Other perennial grasses | 1 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | |
| s5853: | | | | | | <i>Pct</i> |
| Coit | --- | 3,000 | 2,400 | 1,800 | Creeping wildrye | 35 |
| | | | | | Basin wildrye | 15 |
| | | | | | Coyote willow | 10 |
| | | | | | Other perennial grasses | 10 |
| | | | | | Sedge | 10 |
| | | | | | Other perennial forbs | 6 |
| | | | | | Baltic rush | 5 |
| | | | | | Woods rose | 3 |
| | | | | | Cinquefoil | 2 |
| | | | | | Other shrubs | 2 |
| | | | | | Yarrow | 2 |
| Rixie | --- | 3,500 | 2,200 | 1,500 | Basin wildrye | 40 |
| | | | | | Creeping wildrye | 30 |
| | | | | | Other perennial forbs | 10 |
| | | | | | Willow | 5 |
| Rixie | --- | 1,700 | 1,150 | 600 | Basin wildrye | 40 |
| | | | | | Alkali sacaton | 20 |
| | | | | | Inland saltgrass | 15 |
| | | | | | Black greasewood | 10 |
| | | | | | Rubber rabbitbrush | 5 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | <i>Pct</i> |
| s5853: | | | | | | |
| Griver | --- | 3,000 | 2,400 | 1,800 | Basin wildrye | 30 |
| | | | | | Creeping wildrye | 30 |
| | | | | | Baltic rush | 5 |
| | | | | | Coyote willow | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Other shrubs | 5 |
| | | | | | Sedge | 5 |
| | | | | | Woods rose | 5 |
| | | | | | Cinquefoil | 2 |
| | | | | | Groundsel | 2 |
| | | | | | Other perennial forbs | 2 |
| | | | | | Wildiris | 2 |
| | | | | | Yarrow | 2 |
| Griver | --- | 2,000 | 1,500 | 1,000 | Tufted hairgrass | 40 |
| | | | | | Big bluegrass | 10 |
| | | | | | Sedge | 10 |
| | | | | | Other perennial forbs | 8 |
| | | | | | Cinquefoil | 5 |
| | | | | | Meadow barley | 5 |
| | | | | | Other perennial grasses | 5 |
| | | | | | Mannagrass | 3 |
| | | | | | Yarrow | 3 |
| | | | | | Baltic rush | 2 |
| | | | | | Coyote willow | 2 |
| | | | | | Groundsel | 2 |
| | | | | | Wildiris | 2 |
| | | | | | Woods rose | 2 |
| | | | | | Other shrubs | 1 |

Rangeland Productivity and Plant Composition

United States

| Map symbol and soil name | Ecological site | Total dry-weight production | | | Characteristic vegetation | Rangeland composition |
|-----------------------------|-----------------|-----------------------------|----------------|---------------------|---------------------------|--------------------------|
| | | Favorable year | Normal year | Unfavorable year | | |
| | | <i>Lb/ac</i> | <i>Lb/ac</i> | <i>Lb/ac</i> | | |
| s5853: | | | | | | <i>Pct</i> |
| Ocala | --- | 1,900 | 1,400 | 800 | Basin wildrye | 50 |
| | | | | | Alkali sacaton | 20 |
| | | | | | Black greasewood | 10 |
| | | | | | Inland saltgrass | 10 |
| | | | | | Rubber rabbitbrush | 5 |
| Rose Creek | --- | 3,000 | 2,500 | 1,800 | Creeping wildrye | 30 |
| | | | | | Basin wildrye | 25 |
| | | | | | Willow | 15 |
| | | | | | Rush | 5 |
| | | | | | Sedge | 5 |
| | | | | | Woods rose | 5 |
| Rose Creek | --- | 2,500 | 1,900 | 1,200 | Basin wildrye | 55 |
| | | | | | Basin big sagebrush | 10 |
| | | | | | Nevada bluegrass | 10 |
| | | | | | Mat muhly | 5 |
| | | | | | Sedge | 5 |