

# Rangeland Productivity and Plant Composition

United States

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland composition
		Favorable year	Normal year	Unfavorable year		
		<i>Lb/ac</i>	<i>Lb/ac</i>	<i>Lb/ac</i>		<i>Pct</i>
s6501: Wingville	MEADOW	5,000	4,000	3,000	Tufted hairgrass Sedge Rush Bluegrass	65 20 8 5
Baldock	WET MEADOW	3,000	2,500	2,000	Sedge Tufted hairgrass Rush	80 8 5
Haines	SODIC MEADOW	2,500	1,500	1,000	Inland saltgrass Lemmon's alkaligrass Alkali sacaton Rush Alkali cordgrass Basin wildrye	30 30 10 8 5 5
Umapine	SODIC BOTTOM	5,000	4,000	3,000	Basin wildrye Inland saltgrass Black greasewood	80 10 5
Burkemont	SODIC MEADOW	2,500	1,500	1,000	Inland saltgrass Lemmon's alkaligrass Alkali sacaton Rush Alkali cordgrass Basin wildrye	30 30 10 8 5 5

# Rangeland Productivity and Plant Composition

United States

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland composition
		Favorable year	Normal year	Unfavorable year		
		<i>Lb/ac</i>	<i>Lb/ac</i>	<i>Lb/ac</i>		<i>Pct</i>
s6501: Balm	MEADOW	5,000	4,000	3,000	Tufted hairgrass Sedge Rush Bluegrass	65 20 8 5
Stanflow	SODIC BOTTOM	5,000	4,000	3,000	Basin wildrye Inland saltgrass Black greasewood	80 10 5
Boyce	WET MEADOW	3,000	2,000	1,500	Sedge Baltic rush	70 10