## **United States**


[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland
		Favorable year	Normal year	Unfavorable year	Singulation vogetation	composition
		Lb/ac	Lb/ac	Lb/ac		Pct
s9061:						
Paulson	LOAMY (15-19W)	2,400	2,000	1,400	Idaho fescue	20
					Streambank wheatgrass	15
					Antelope bitterbrush	10
					Big sagebrush	10
					Canby bluegrass	10
					Spike fescue	10
					California brome	5
					Columbia needlegrass	5
Leavittville	LOAMY (15-19W)	2,400	2,000	1,400	Idaho fescue	20
					Streambank wheatgrass	15
					Antelope bitterbrush	10
					Big sagebrush	10
					Canby bluegrass	10
					Other perennial forbs	10
					Spike fescue	10
					California brome	5
					Columbia needlegrass	5
					Other perennial grasses	5



## **United States**

Map symbol and soil name	Ecological site	Total o	lry-weight prod	uction	Characteristic vegetation	Rangeland
		Favorable year	Normal year	Unfavorable year		composition
		Lb/ac	Lb/ac	Lb/ac		Pc
s9061:						
Buckskin	LOAMY (15-19W)	2,400	2,000	1,400	Bluebunch wheatgrass	30
					Mountain big sagebrush	10
					Other perennial grasses	10
					Antelope bitterbrush	5
					Arrowleaf balsamroot	5
					Columbia needlegrass	5
					Cutleaf balsamroot	5
					Geranium	5
					Mountain brome	5
					Nevada bluegrass	5
					Other perennial forbs	5
					Other shrubs	5
					Prairie Junegrass	5
					Slender wheatgrass	5
Hobacker	LOAMY (15-19W)	2,400	2,000	1,400	Bluebunch wheatgrass	20
	, ,				Idaho fescue	20
					Big sagebrush	10
					Antelope bitterbrush	5
					Basin wildrye	5
					Mountain brome	5
					Sandberg bluegrass	5
Decross	LOAMY (15-19W)	2,400	2,000	1,400	Idaho fescue	20
		_,	_,	1,100	Streambank wheatgrass	15
					Antelope bitterbrush	10
					Big sagebrush	10
					Canby bluegrass	10
					Spike fescue	10
					Columbia needlegrass	5



## **United States**

Map symbol and soil name	Ecological site	Total o	dry-weight prod	uction	Characteristic vegetation	Rangeland
		Favorable year	Normal year	Unfavorable year		composition
	•	Lb/ac	Lb/ac	Lb/ac		Pct
s9061:						
Cowdrey						
Robana	LOAMY (15-19W)	2,400	2,000	1,400	Idaho fescue	20
rtobana	EG/WIT (10 10VV)	2,400	2,000	1,400	Streambank wheatgrass	15
					Antelope bitterbrush	10
					Big sagebrush	10
					Canby bluegrass	10
					Other perennial forbs	10
					Spike fescue	10
					California brome	5
					Columbia needlegrass	5
					Other perennial grasses	5
Greyback	LOAMY (15-19W)	2,400	2,000	1,400	Idaho fescue	20
					Streambank wheatgrass	15
					Antelope bitterbrush	10
					Big sagebrush	10
					Canby bluegrass	10
					Other perennial forbs	10
					Spike fescue	10
					California brome	5
					Columbia needlegrass	5
					Other perennial grasses	5



## **United States**

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland
		Favorable year	Normal year	Unfavorable year	Onardolonollo Vogolallon	composition
		Lb/ac	Lb/ac	Lb/ac		Pct
s9061:						
Turson	SUBIRRIGATED (15-19W)	5,500	4,500	3,500	Tufted hairgrass	20
					Kentucky bluegrass	10
					Other perennial forbs	10
					Slender wheatgrass	10
					Willow	10
					Alpine timothy	5
					Basin wildrye	5
					California brome	5
					Nebraska sedge	5
					Other perennial grasses	5
					Sedge	5
					Shrubby cinquefoil	5
					Western wheatgrass	5

